

Noumed Pioglitazone

pioglitazone hydrochloride tablets

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Noumed Pioglitazone. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Noumed Pioglitazone against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What this medicine is used for

Noumed Pioglitazone helps to control the level of glucose in your blood when you have type 2 diabetes. This is the 'adult onset' type of diabetes and is controlled by diet, certain oral medications and occasionally insulin.

This medicine is used:

- to improve the action of the body's naturally produced insulin.
- in the management of type 2 diabetes not controlled by diet.
- alone (when diet and exercise is not enough to treat your diabetes) or together with other anti-diabetic medicines.

It contains the active ingredient pioglitazone hydrochloride.

Pioglitazone hydrochloride belongs to a group of medicines called glitazones.

It works by decreasing insulin resistance.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

This medicine is not addictive.

This medicine is available only with a doctor's prescription.

The use of Noumed Pioglitazone has not been studied in children.

Before you take this medicine

When you must not take it

Do not take this medicine if you have an allergy to:

- Pioglitazone hydrochloride, the active ingredient, or to any of the other ingredients listed at the end of this leaflet under Product Description.
- any other similar medicines (Rosiglitazone maleate).

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

Do not take this medicine if you have or have had any of the following medical conditions:

- heart failure requiring treatment. Talk to your doctor if you have heart failure
- type 1 diabetes or diabetic ketoacidosis
- bladder cancer or the presence of blood in your urine.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- heart disease with shortness of breath after minimal physical activity
- heart disease with severe symptoms at rest
- swelling of hands, ankles or feet
- bladder cancer or symptoms associated with bladder cancer such as blood in the urine (hematuria) often accompanied by pain and burning or sudden urges to urinate.
- problems with your liver
- problems with your kidneys that requires dialysis. Noumed

Pioglitazone is not recommended for use if you are on dialysis

- some women who do not have monthly periods and have not been through menopause may restart their periods when taking Noumed Pioglitazone. These women may be at increased risk of pregnancy
- bone fractures, usually in the hand, upper arm or foot, have been seen in some women when taking Noumed Pioglitazone. Talk to your doctor for advice on how to keep your bones healthy.

Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.

Your doctor can discuss with you the risks and benefits involved.

Tell your doctor if you are using another medicine for diabetes. Noumed Pioglitazone can enhance the action of other medicines. You may be at risk of low blood sugar (hypoglycaemia).

If this happens, your doctor may need to adjust the dose of your other medicines.

If you have not told your doctor about any of the above, tell him/her before you start taking Noumed Pioglitazone tablets.

Taking other medicines

Tell your doctor if you are taking any other medicine, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Noumed Pioglitazone may interfere with each other. These include:

- gemfibrozil, a medicine used to lower high cholesterol
- other antidiabetic drugs such as glibenclamide, gliclazide, insulin, metformin, tolbutamide and chlorpropamide
- oral contraceptives
- rifampicin, an antibiotic drug.

These medicines may be affected by Noumed Pioglitazone or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to take this medicine

Follow all directions given to you by your doctor or pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box/bottle, ask your doctor or pharmacist for help.

How much to take

Your doctor will tell you how many Noumed Pioglitazone tablets you should take.

The standard dose for this medicine is usually in the range of 15 mg to 45 mg per day.

Your doctor may increase or decrease your dose in order to find the appropriate dose for your condition. How to take it

Swallow the tablet whole with a full glass of water.

When to take it

Noumed Pioglitazone tablets should be taken once a day.

Take your medicine at about the same time each day.

Taking it at the same time each day will have the best effect. It will also help you remember when to take it.

It does not matter whether you take Noumed Pioglitazone with food or on an empty stomach.

How long to take it

Continue taking your medicine for as long as your doctor tells you.

This medicine helps to control your condition, but does not cure it. It is important to keep taking your medicine even if you feel well.

If you forget to take it

Take your dose as soon as you remember, and continue to take it as you would normally.

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much [Medicine name]. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

While you are using this medicine

Things you must do

It is important that you remember to take Noumed Pioglitazone daily

and at the dose prescribed by your doctor.

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking Noumed Pioglitazone.

Tell any other doctors, dentists and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.

It may affect other medicines used during surgery.

If you become pregnant while taking this medicine, tell your doctor immediately.

If you are about to have any blood tests, tell your doctor that you are taking this medicine.

It may interfere with the results of some tests.

Keep all of your doctor's appointments so that your progress can be checked.

Your doctor may do some tests from time to time to make sure the medicine is working and to prevent unwanted side effects.

Tell your doctor if you have gained weight since taking Noumed Pioglitazone.

Weight gain can be associated with improved blood sugar control however; it may also be a symptom of heart failure.

Things you must not do

Do not take Noumed Pioglitazone to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your medicine or lower the dosage without checking with your doctor.

If you stop taking it suddenly, your condition may worsen or you may have unwanted side effects.

Things to be careful of

Noumed Pioglitazone alone is unlikely to affect your ability to drive or operate machinery. However, be careful to avoid hypoglycaemia whilst driving or operating machinery if using Noumed Pioglitazone in combination with other diabetes medicines.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Noumed Pioglitazone.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

This medicine helps most people with type 2 diabetes not controlled by diet, but it may have unwanted side effects in some people.

Some side effects may be related to the dose of Noumed Pioglitazone. Accordingly, it is important that you tell your doctor as soon as possible about any unwanted effects. Your doctor may then decide to adjust the dose of Noumed Pioglitazone you are taking.

Do not be alarmed by the following list of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- weight gain
- signs of low blood sugar levels (hypoglycaemia) which may include weakness, trembling or shaking, sweating, light-headedness, headache, dizziness, lack of concentration, tearfulness or crying, irritability, hunger,

numbness around the lips and fingers

- Eye problems including blurred or double vision.

These are more common side effects of the medicine.

Tell your doctor as soon as possible if you notice any of the following:

- dark urine or pale stools, yellowing of the skin or eyes, severe cramps of the stomach, nausea or vomiting, loss of weight, tiredness
- shortness of breath when at rest or after minimal physical activity with swelling of legs, feet and hands, rapid increase in weight
- blood in the urine often accompanied by pain and burning, these can be symptoms of bladder cancer.

The above list includes serious side effects that may require medical attention. Serious side effects are rare.

Tell your doctor or pharmacist if you notice anything else that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After using this medicine

Storage

Keep your medicine in the original container.

If you take it out of its original container it may not keep well.

Keep your medicine in a cool dry place where the temperature stays below 30°C.

Do not store Noumed Pioglitazone or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car.

Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Level 5, 7 Eden Park Drive,
Macquarie Park, NSW 2113
Australia
Phone: 1800 930 999

This leaflet was prepared in October 2020.

Australian Register Number:

15mg tablets: AUST R 285673

30mg tablets: AUST R 285674

45mg tablets: AUST R 285675

Product description

What it looks like

Noumed Pioglitazone comes in three tablet strengths:

Noumed Pioglitazone 15mg - white, round tablet, with imprint "PGT 15" on one side.

Noumed Pioglitazone 30mg - white, round tablet, with imprint "PGT 30" on one side and with score line on both sides.

Noumed Pioglitazone 45mg - white, round tablet, with imprint "PGT 45" on one side.

Available in blister packs of 28 tablets.

Ingredients

Noumed Pioglitazone contains 15 mg, 30 mg or 45 mg of pioglitazone (as hydrochloride) as the active ingredient.

It also contains the following inactive ingredients:

- lactose monohydrate
- hypolose
- carmellose calcium
- magnesium stearate

This medicine does not contain gluten, tartrazine or any other azo dyes.

Sponsor

Avallon Pharmaceuticals Pty Ltd